

This is a repository copy of *Psychophysical measurement of the effects and non-effects of TMS on contrast perception*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/129411/>

Version: Accepted Version

Article:

Vilidaite, Greta and Baker, Daniel Hart orcid.org/0000-0002-0161-443X (2018)
Psychophysical measurement of the effects and non-effects of TMS on contrast perception. *Brain Stimulation*. pp. 956-957. ISSN 1935-861X

<https://doi.org/10.1016/j.brs.2018.04.005>

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Accepted Manuscript

Psychophysical measurement of the effects and non-effects of TMS on contrast perception

Greta Vilidaite, Daniel H. Baker

PII: S1935-861X(18)30106-2

DOI: [10.1016/j.brs.2018.04.005](https://doi.org/10.1016/j.brs.2018.04.005)

Reference: BRS 1231

To appear in: *Brain Stimulation*


Received Date: 2 March 2018

Revised Date: 28 March 2018

Accepted Date: 4 April 2018

Please cite this article as: Vilidaite G, Baker DH, Psychophysical measurement of the effects and non-effects of TMS on contrast perception, *Brain Stimulation* (2018), doi: 10.1016/j.brs.2018.04.005.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



1 **Title: Psychophysical measurement of the effects and**
2 **non-effects of TMS on contrast perception**

3

4

5

6 **Greta Vilidaite**, Department of Psychology, University of York, York, YO10
7 5DD, United Kingdom

8

9 **Daniel H Baker***, Department of Psychology, University of York, York, YO10
10 5DD, United Kingdom

11

12

13

14

15

16

17 * Corresponding author. Email: daniel.baker@york.ac.uk

18 **Highlights**

19

- 20 • Compared the effects of four TMS protocols on neural signals and
21 noise.
- 22 • Single pulse TMS suppressed neural signals and repetitive TMS
23 increased neural noise.
- 24 • Theta burst stimulation did not affect perceptual task performance.
- 25 • Participants differed in TMS susceptibility, determined by phosphene
26 perception.
- 27 • Findings suggest systematic inter-protocol and inter-participant
28 differences in TMS effects.

29

30

31 **Key words**

32 Sensory processing, neural effects, theta burst, online stimulation,
33 psychophysics

To the editor:

Transcranial magnetic stimulation (TMS) is widely used to establish causal relationships between brain areas and behavior, but its effects on task performance are not fully understood and have rarely been directly compared between protocols. Decreases in performance on psychophysical tasks, such as those observed when applying TMS, can be attributed to either suppression of stimulus-related neural signals, increased random activity (i.e. neural noise), or a combination of both [1,2]. Indeed, evidence for all three hypotheses has been found when using differing methodologies and online stimulation protocols [3–5]. Similarly, theta burst stimulation (TBS) has been shown to have variable or bimodal effects between participants and between exact stimulation protocols [6,7]. Despite different TMS protocols (e.g. online, offline, repetitive, single pulse) potentially having vastly different effects, they are often used interchangeably in sensory and cognitive research.

We directly compared the neural effects of four commonly used TMS protocols: online single pulse (spTMS), online 3-pulse repetitive (rTMS; 50ms between pulses), offline continuous theta burst (cTBS) and offline intermittent theta burst (iTBS), during a well-understood neural computation – contrast transduction. As a secondary objective, we investigated natural TMS-susceptibility by comparing participants who could and could not perceive phosphenes to address inter-participant variability in TMS effectiveness.

We tested all stimulation protocols using the same area (occipital cortex, Supp. 2A) and a highly sensitive double-pass paradigm [8] to dissociate TMS induced changes in stimulus-related neural signal strength (i.e. suppression) and neural noise. On each trial (200 total per TMS condition) two luminance-modulated stimuli (3 deg. vis. ang.) of randomly-selected contrast were presented peripherally. Half of the trials contained a 4% contrast increment in one of the intervals (see Supp. 1A,B for examples). The exact same trials were then repeated with randomized interval order. Full details of stimuli and the double-pass paradigm be found in [9]. Using standard protocols with a Magstim Super Rapid² 'figure of 8' coil spTMS and rTMS (Supp. 2C, 70% stimulator output) were applied 50ms after stimulus onset in each interval, and offline TBS (Supp. 2D, 30% stimulator output) was applied before the start of the task. Consistency between the first and second presentation of the trials was calculated as a direct index of neural noise. Accuracy on the task was calculated as a measure of stimulus-related signal strength.

During phosphene localization pre-screening, six participants (4 females, age 22-34) consistently perceived phosphenes and completed the main experiment (a further 19 participants were screened but did not report seeing phosphenes). Study was approved by YNiC ethics committee. All TMS protocols were tested on different days (rTMS was tested over four days due to high numbers of pulses). Phosphene localization was performed before each testing session and the location of the phosphenes (as indicated with a computer interface, Fig. 1A) was used to subsequently present stimuli.

Control trials (no TMS) were performed before stimulation for each TMS protocol separately.

We simulated predictions using a linear amplifier model (LAM). Simulations showed that if TMS reduced neural signal strength (lowered sensitivity), we would observe a steep drop in task accuracy but no change in double-pass consistency. Alternatively, if TMS increased neural noise, we would see a small reduction in accuracy and a larger drop in consistency. Finally, if TMS both reduced stimulus-related signals and increased noise, we would observe a large reduction in both measures (Supp. 1C-E).

We found a significant drop in accuracy ($t(5)=2.83$, $p=0.037$, Bayes factor (BF)=2.83) when applying spTMS compared to the no TMS condition, but no change in consistency ($p=0.601$, BF=0.29, Fig. 1B). This closely resembles our LAM model predictions for an increase in neural suppression and suggests that spTMS suppresses neural signals. Conversely, applying rTMS showed a small non-significant change in accuracy ($p=0.848$, BF=0.33) compared to the no-TMS condition, and a significant decrease in consistency ($t(5)=2.74$, $p=0.041$, BF=2.38, Fig. 1C) – consistent with model predictions for an increase in neural noise. Neither protocol produced data consistent with change in both suppression and noise. This comparison between spTMS and rTMS is consistent with previous research that tested these protocols separately [4,3] and suggests suppressive and noise-inducing effects are protocol-specific.

No effects on the accuracy ($p=0.790$, $BF=0.30$) or consistency ($p=0.132$, $BF=0.93$) were observed when applying cTBS (Fig. 1D). Similarly, no changes in accuracy ($p=0.773$, $BF=0.30$) or consistency ($p=0.244$, $BF=0.58$) were observed when applying iTBS (Fig. 1E), indicating that neither protocol changed the levels of neural noise or sensory signals. This may seem to oppose the large number of successful TBS studies, particularly in the motor cortex. However, most previous research into TBS effects measured motor evoked potentials, which reflect an overall increase or decrease in neural activity (e.g. [10]). It may be that TBS changes overall neural activation but does not have particular effects on perceptually-relevant signals that would affect sensory task performance. Alternatively, the effectiveness of TBS may be overstated in the literature, as indicated by a recent large scale meta-analysis [11] which found a large positive publication bias in the TBS literature.

To investigate the effects of TMS susceptibility on task-relevant effects, a further six participants (3 females, age 23-55) who did not report seeing phosphenes also completed the experiment. For these participants, stimuli were presented at the mean location of phosphenes experienced by the other group. None of the four TMS protocols had any significant effect on accuracy or consistency scores in these individuals, indicating that the participants who did not perceive phosphenes during phosphene localization were not affected by TMS during the task. Anatomical differences in cortical folding and skull thickness may explain these individual differences in TMS susceptibility.

The inter-participant and inter-protocol differences in TMS effects found here shed light on the interpretation of findings in the existing TMS literature and inform future methodological choices. The individual differences in susceptibility and the use of different stimulation protocols in the literature may be some of the major factors in the TMS ‘replication crisis’ [12]. The effects of TMS are subtle and can often only be detected in reaction time data rather than task performance [13]. In this respect, the sensitivity and precision of the double-pass paradigm is a valuable tool for further investigating TMS inter-protocol and inter-participant variability in other brain areas and with larger samples.

Conflict of Interest

There is no conflict of interest relating to this manuscript.

References

- [1] Walsh V, Cowey A. Magnetic stimulation studies of visual cognition. *Trends Cogn Sci* 1998;2:103–10. doi:10.1016/S1364-6613(98)01134-6.
- [2] Walsh V, Pascual-Leone A. Transcranial magnetic stimulation: a neurochronometrics of mind. Cambridge, Mass; London: MIT Press; 2003.
- [3] Harris JA, Clifford CWG, Miniussi C. The functional effect of transcranial magnetic stimulation: signal suppression or neural noise generation? *J Cogn Neurosci* 2008;20:734–40. doi:10.1162/jocn.2008.20048.
- [4] Ruzzoli M, Abrahamyan A, Clifford CWG, Marzi CA, Miniussi C, Harris JA. The effect of TMS on visual motion sensitivity: an increase in neural

- 157 noise or a decrease in signal strength? J Neurophysiol 2011;106:138–
 158 43. doi:10.1152/jn.00746.2010.
- 159 [5] Moliadze V, Zhao Y, Eysel U, Funke K. Effect of transcranial magnetic
 160 stimulation on single-unit activity in the cat primary visual cortex. J
 161 Physiol 2003;553:665–79. doi:10.1113/jphysiol.2003.050153.
- 162 [6] Harada M, Taki MM, Naa MÁ. Non-Invasive Evaluation of the
 163 GABAergic / Glutamatergic System in Autistic Patients Observed by
 164 MEGA-Editing Proton MR Spectroscopy Using a Clinical 3 Tesla
 165 Instrument. J Autism Dev Disord 2011;41:447–54. doi:10.1007/s10803-
 166 010-1065-0.
- 167 [7] Gentner R, Wankerl K, Reinsberger C, Zeller D, Classen J. Depression
 168 of human corticospinal excitability induced by magnetic theta-burst
 169 stimulation: evidence of rapid polarity-reversing metaplasticity. Cereb
 170 Cortex 2008;18:2046–53. doi:10.1093/cercor/bhm239.
- 171 [8] Burgess AE, Colborne B. Visual signal detection. IV. Observer
 172 inconsistency. J Opt Soc Am 1988;5:617–27.
 173 doi:10.1364/JOSAA.5.000617.
- 174 [9] Vilidaite G, Yu M, Baker DH. Internal noise estimates correlate with
 175 autistic traits. Autism Res 2017;10:1384–1391. doi:10.1002/aur.1781.
- 176 [10] Di Lazzaro V, Pilato F, Dileone M, Profice P, Oliviero A, Mazzone P, et
 177 al. The physiological basis of the effects of intermittent theta burst
 178 stimulation of the human motor cortex. J Physiol 2008;586:3871–9.
 179 doi:DOI 10.1113/jphysiol.2008.152736.
- 180 [11] Chung SW, Hill AT, Rogasch NC, Hoy KE, Fitzgerald PB. Use of theta-
 181 burst stimulation in changing excitability of motor cortex: A systematic

- 182 review and meta-analysis. *Neurosci Biobehav Rev* 2016;63:43–64.
183 doi:10.1016/j.neubiorev.2016.01.008.
- 184 [12] Héroux ME, Taylor JL, Gandevia SC. The use and abuse of transcranial
185 magnetic stimulation to modulate corticospinal excitability in humans.
186 *PLoS One* 2015;10:1–10. doi:10.1371/journal.pone.0144151.
- 187 [13] Pitcher D. Facial Expression Recognition Takes Longer in the Posterior
188 Superior Temporal Sulcus than in the Occipital Face Area
189 2014;34:9173–7. doi:10.1523/JNEUROSCI.5038-13.2014.
190

Figures






Figure 1. Phosphene locations, mean accuracy and consistency scores for the individuals seeing phosphenes. Phosphene locations were similar for all six participants, centered around the midline of the left visual field (A), within 15 degrees of the fixation cross. Phosphene locations were consistent across the four experiments using different stimulation protocols: spTMS (blue), rTMS (yellow,

199 averaged over four sessions), cTBS (green) and iTBS (purple), as indicated by filled
200 ovals. In Exp 1, single pulse TMS (B) significantly reduced the mean accuracy scores
201 (dark bars) compared to the no-TMS condition (light bars) but not consistency scores
202 which indicates increased suppression resulting from TMS stimulation. Repetitive
203 TMS (C) significantly reduced task consistency but not task accuracy, indicating a
204 TMS-induced increase in neural noise. Neither cTBS (D) nor iTBS (E) produced any
205 significant change in task performance. Error bars indicate bootstrapped 95%
206 confidence intervals.

Supplementary materials



Supplementary figure 1. Each interval during a trial was drawn from the target (blue) and non-target (yellow) stimulus distributions (A). Participants were asked to choose the interval with the more positive contrast (B; example correct intervals are shown with a blue circle). Stochastic simulations were used to generate model predictions of double-pass data (C-E). Light bars in all panels indicate a system with low neural noise and low suppression (high sensitivity) in the system. Dark bars model an increase in either suppression, noise, or both. If TMS suppresses neural signals (lowers sensitivity) then we should expect double-pass data to be similar to the prediction in panel C. On the other hand, if TMS increases neural noise the data should resemble panel D. If both suppression and neural noise are increased we would expect data to be similar to panel E.



221

222 **Supplementary figure 2.** The TMS coil was positioned (red dot) approximately 2cm
 223 above and 1cm to the right of the inion (blue line intersection) to induce phosphenes
 224 (A). Before phosphene localization participants were trained to indicate the location
 225 and shape of a simulated phosphene on the screen (B; see section 2.3). During
 226 spTMS and rTMS protocols either one or three pulses (50ms apart) were delivered
 227 50ms after stimulus onset (C). Pulses during offline cTBS and iTBS were delivered
 228 as shown in D.